

# Prostaglandin E(2) induces spontaneous rhythmic activity in mouse urinary bladder independently of efferent nerves

Kobayter, S; Young, JS; Brain, Keith

DOI:

[10.1111/j.1476-5381.2011.01543.x](https://doi.org/10.1111/j.1476-5381.2011.01543.x)

## Document Version

Publisher's PDF, also known as Version of record

## Citation for published version (Harvard):

Kobayter, S, Young, JS & Brain, K 2012, 'Prostaglandin E(2) induces spontaneous rhythmic activity in mouse urinary bladder independently of efferent nerves', *British Journal of Pharmacology*, vol. 165, no. 2, pp. 401-413. <https://doi.org/10.1111/j.1476-5381.2011.01543.x>

[Link to publication on Research at Birmingham portal](#)

## General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

## RESEARCH PAPER

# Prostaglandin E<sub>2</sub> induces spontaneous rhythmic activity in mouse urinary bladder independently of efferent nerves

S Kobayter<sup>1</sup>, JS Young<sup>1\*</sup> and KL Brain<sup>2</sup>

<sup>1</sup>Department of Pharmacology, University of Oxford, Oxford, UK, and <sup>2</sup>School of Clinical and Experimental Medicine, University of Birmingham, Medical School, Birmingham, UK

### Correspondence

KL Brain, School of Clinical and Experimental Medicine, University of Birmingham, IBR Wolfson Drive, Medical School, Birmingham B15 2TT, UK. E-mail: k.l.brain@bham.ac.uk

\*Present address: Faculty of Health and Medical Sciences, Wolfson Cell Science Building, University of Surrey, Guildford, Surrey GU2 7XH, UK.

Re-use of this article is permitted in accordance with the Terms and Conditions set out at [http://wileyonlinelibrary.com/onlineopen#OnlineOpen\\_Terms](http://wileyonlinelibrary.com/onlineopen#OnlineOpen_Terms)

### Keywords

prostaglandin E; urinary bladder; smooth muscle; calcium; parasympathetic nervous system

### Received

26 July 2010

### Revised

19 April 2011

### Accepted

1 June 2011

## BACKGROUND AND PURPOSE

The acute effects of PGE<sub>2</sub> on bladder smooth muscle and nerves were examined to determine the origin of PGE<sub>2</sub>-induced spontaneous rhythmic contractions.

## EXPERIMENTAL APPROACH

Contraction studies, confocal Ca<sup>2+</sup> imaging and electrophysiological recordings in strips of mouse urinary bladder were used to differentiate the effects of PGE<sub>2</sub> on bladder smooth muscle and efferent nerves.

## KEY RESULTS

PGE<sub>2</sub> (50 µM) increased the tone and caused phasic contractions of detrusor smooth muscle strips. Confocal Ca<sup>2+</sup> imaging showed that PGE<sub>2</sub> increased the frequency of whole-cell Ca<sup>2+</sup> transients (WCTs) (72 ± 5%) and intracellular recordings showed it increased the frequency of spontaneous depolarizations, from 0.31·s<sup>-1</sup> to 0.90·s<sup>-1</sup>. Non-selective inhibition of EP receptors using SC-51322 and AH-6809 (10 µM), or the L-type Ca<sup>2+</sup> channel blocker nifedipine (1 µM), prevented these phasic contractions and WCTs, and reduced the tone (by 45 ± 7% and 59 ± 6%, respectively). Blocking P2X1 receptors with NF449 (10 µM) caused a small but significant reduction in the frequency of PGE<sub>2</sub>-induced phasic contractions (24 ± 9%) and WCTs (28 ± 17%) but had no significant effect on spontaneous depolarizations or tone. Inhibiting muscarinic receptors with cyclopentolate (1 µM) had no significant effect on these measures. Spontaneous WCTs became synchronous in PGE<sub>2</sub>, implying enhanced functional coupling between neighbouring cells. However, the electrical input resistance was unchanged.

## CONCLUSIONS AND IMPLICATIONS

It was concluded that depolarization alone is sufficient to explain a functional increase in intercellular coupling and the ability of PGE<sub>2</sub> to increase detrusor spontaneous rhythmic activity does not require parasympathetic nerves.

## Abbreviations

AH-6809, 6-isopropoxy-9-xanthone-2-carboxylic acid; KO, knockout; NF449, 4,4',4'',4'''-[carbonylbis(imino-5,1,3-benzenetriyl-bis(carbonylimino))]tetrakis-1,3-benzenedisulphonic acid; sAP, spontaneous action potential; sEJP, spontaneous excitatory junction potential; SC-51322, 8-chloro-2-[3-[(2-furanylmethyl)thio]-1-oxopropyl]-dibenz(Z)[b,f][1,4]oxazepine-10(11H)-carboxylic acid hydrazide; SMC, smooth muscle cell; TTX, tetrodotoxin; WCT, whole-cell Ca<sup>2+</sup> transient

## Introduction

Prostanoids are potent and ubiquitous inflammatory mediators. In the bladder, production of prostanoids, particularly PGE<sub>2</sub>, is elevated in pathological conditions, such as detrusor overactivity that follows spinal cord injury (Masunaga *et al.*, 2006), and bladder outlet obstruction (Schröder *et al.*, 2004). It is likely that this association is at least partially causative, because inhibiting COX with indomethacin reduces the symptoms of bladder overactivity in people with neurogenic detrusor instability (Cardozo and Stanton, 1980). Prostanoid synthesis occurs locally in bladder urothelium and smooth muscle. This synthesis can be initiated by factors including stretch, nerve stimulation, injury, exposure to ATP and other inflammatory mediators (Dveksler *et al.*, 1989; Khan *et al.*, 1998).

Prostanoids are implicated in the modulation of bladder function (Khan *et al.*, 1998). If PGE<sub>2</sub> is instilled intravesically in the conscious, catheterized rat, micturition is facilitated and an increased basal intravesical pressure is generated (Ishizuka *et al.*, 1995). Similar urodynamic effects are observed in humans; intravesical instillation of PGE<sub>2</sub> causes detrusor overactivity, urgency and decreased bladder capacity (Schussler, 1990). The basis for such excitatory effects, observed in detrusor strips of animal models and humans (Maggi, 1992) is likely to be a slow increase in detrusor muscle tone, followed by an increase in phasic contractions (Bultitude *et al.*, 1976). These PG-generated smooth muscle contractions are inhibited by the L-type calcium channel blocker verapamil (Hanawa, 1991; cited in Khan *et al.*, 1998).

PGE<sub>2</sub> is likely to mediate its effects through more than one EP receptor. The competitive EP<sub>1</sub> antagonist SC-19220 increases bladder capacity and reduces voiding efficiency in anaesthetized rats (Maggi *et al.*, 1988). Recent studies in EP<sub>1</sub> knockout (KO) mice have shown that PGE<sub>2</sub> is not essential for normal micturition but does play a role in detrusor overactivity (Schröder *et al.*, 2004). Studies using EP receptor KO mice suggest a role for EP<sub>1</sub> in detrusor overactivity (Schröder *et al.*, 2004), whereas EP<sub>3</sub> is implicated in bladder capacity in physiology (McCafferty *et al.*, 2008). Further experiments are required to characterize the role of different EP receptors in physiological and pathophysiological conditions.

It is still unclear whether PGE<sub>2</sub> mediates its actions in the bladder directly on smooth muscle cells (SMCs) or indirectly by acting on nerves. It has been suggested that PGs released from SMCs and from the urothelium modulate afferent nerve activity and efferent neurotransmission (de Groat and Yoshimura, 2001). Furthermore, our group recently showed that spontaneous action potentials (sAPs) in mouse bladder are purinergic and neurogenic (Young *et al.*, 2008), suggesting that if PGE<sub>2</sub> affects spontaneous activity it could do so through a direct action on efferent nerves.

Hence, the aim of this study was to differentiate the effects of PGE<sub>2</sub> on bladder smooth muscle and efferent nerves using a combination of contraction studies, high-resolution Ca<sup>2+</sup>-imaging and electrophysiology.

## Methods

Female CD-1 mice, aged 8–13 weeks, were killed by increasing the CO<sub>2</sub> concentration followed by cervical dislocation. All experiments were carried out in accordance with the UK Animals (Scientific Procedures) Act 1986 and European Communities Council Directive 86/09/EEC. The urinary bladder was removed and placed in physiological saline. The bathing physiological salt solution contained (mM): NaCl 118.4, NaHCO<sub>3</sub> 25.0, NaH<sub>2</sub>PO<sub>4</sub> 1.13, CaCl<sub>2</sub> 1.8, KCl 4.7, MgCl<sub>2</sub> 1.3 and glucose 11.1. pH and [O<sub>2</sub>] were regulated by continuously bubbling the solution reservoir with 95% O<sub>2</sub> and 5% CO<sub>2</sub>. The connective tissue surrounding the bladder was first removed, then the ventral wall of the bladder was opened longitudinally from the bladder neck to the apex of the dome. The urothelium was left intact. Tissue strips, 6–8 mm long and 1–2 mm wide, were cut along the craniocaudal axis of the dorsal surface for organ bath studies and were 3–4 mm long and 1–2 mm wide for electrophysiological recording or Ca<sup>2+</sup> imaging.

### Contraction studies

Each bladder strip was mounted in a 7 mL organ bath and connected to an isometric transducer (Letica Scientific Instruments, Hospitalet, Spain), under an initial tension of 9.8 mN and allowed to equilibrate (i.e. to accommodate under tension) for at least 60 min. Contraction data were digitized using a Powerlab/4SP data acquisition system, Chart v.5.5.6/s software (ADInstruments, Chalgrove, UK) and data stored on a Macintosh computer. The maximum amplitude of contractions was measured off-line.

### Electrophysiological recordings

Bladder strips were allowed to equilibrate for at least 30 min, and then individual bladder SMCs were impaled from the serosal surface using a shallow angle of approach and superficial penetration with glass capillary microelectrodes filled with 0.5 M KCl (resistance 200–300 MΩ). This approach is identical to that used previously in our laboratory (Young *et al.*, 2008), where filling the cells with Oregon Green 488 BAPTA-1 (mixed with the KCl in the microelectrode) showed large cells typical of SMCs (Dixon and Gosling, 1990; Andersson and Arner, 2004). For both electrophysiology and Ca<sup>2+</sup> imaging, pinning with a small amount of even stretch reduced most spontaneous movement of detrusor strips. The use of long-tipped electrodes assisted the maintenance of recordings during movement of the tissue, a particular methodological challenge when using PGE<sub>2</sub>. Following bridge balancing and neutralization of the electrode's tip capacitance, membrane potential changes were recorded using a high input impedance amplifier (Axoclamp-2B; Axon Instruments, Sunnyvale, CA, USA) and displayed on a digital oscilloscope (DSO 1602; Gould, Ilford, Essex, UK). Membrane potential changes were digitized at 1 kHz using a PowerLab/4SP (ADInstruments, Chalgrove, Oxfordshire, UK) and stored on computer for later analysis.

One long, continuous recording per preparation was made, which included a control period (typically 10 min), and then a period (at least 15 min) in the presence of the drug. The approach of making a single recording per prepa-

ration, rather than sampling several SMCs in control conditions and then several SMCs following perfusion of a drug, was favoured as the PGE<sub>2</sub>-induced contractions greatly hindered subsequent impalement of further SMCs.

The electrical properties of mouse detrusor smooth muscles were determined by passing outward current (in steps of  $-0.03$  to  $-0.15$  nA, of 100 ms duration) through the recording microelectrode.

To determine the frequency with which bursts of spontaneous depolarizations occurred following PGE<sub>2</sub> superfusion, recording segments in which such bursts were present were assessed. The membrane potential was low-pass filtered (0.1 to 0.5 Hz cut-off frequency) and a correlation coefficient of this channel to a typical low-pass filtered burst (template) was calculated. Bursts were then defined using the 'cycle variables' function, with the noise threshold (10 to 40%) adjusted until it provided specificity comparable to manual counting of such bursts in unfiltered data (Chart; ADInstruments, Chalgrove, Oxfordshire, UK).

### Ca<sup>2+</sup> imaging and analysis

Each detrusor strip was exposed to 10  $\mu$ M Oregon Green 488 BAPTA 1-AM (Invitrogen, Paisley, UK) in 1% DMSO-0.2% Pluronic F-127 (Sigma-Aldrich, Dorset, England) in physiological saline for 70 min at 36°C. The tissue was rinsed in physiological saline, bubbled with 95%O<sub>2</sub>/5%CO<sub>2</sub> to pH 7.4, for at least 10 min. Tissues were pinned flat, serosal side up, in a Sylgard-lined organ bath, and mounted on the stage of an upright confocal microscope (Leica SP2; Leica Microsystems, Milton Keynes, UK). A series of 70 frames was captured at 5 Hz using a  $\times 40$  objective to generate one image set. Such sets were acquired once every minute. Five sets were sampled for control and drug treatments within one region, containing a minimum of two cells sampled per drug or treatment. Image analysis was performed using custom-written macros for Image SXM v1.85 (<http://www.liv.ac.uk/~sdb/ImageSXM>). Data were exported to Chart v5.5.6 and then to Excel (Microsoft, Redmond, WA, USA) for further analysis.

### Correlating whole-cell Ca<sup>2+</sup> transients between neighbouring cells

The total number of whole-cell Ca<sup>2+</sup> transients (WCTs) observed in neighbouring cells, before and after superfusion with PGE<sub>2</sub> (50  $\mu$ M) was compared. Only directly neighbouring cells that exhibited WCTs under control conditions were analysed. Contingency analysis (with Fisher's exact test) was used to determine whether cells were functionally coupled.

### Drugs

All drugs were stored at  $-20^{\circ}\text{C}$  as 10 mM stock solutions; PGE<sub>2</sub> and SC-51322 in ethanol, nifedipine and AH-6809 in DMSO, and NF449, tetrodotoxin (TTX) and cyclopentolate in distilled water. Nifedipine and cyclopentolate hydrochloride were obtained from Sigma (Dorset, UK); SC-51322 and AH 6809 were obtained from Biomol (Exeter, UK); and PGE<sub>2</sub>, TTX and NF449 were obtained from Tocris (Bristol, UK). Drugs were diluted to their final concentration with physiological saline and applied directly into the relevant baths in the contraction studies, and by switching the superfusate in the Ca<sup>2+</sup>-imaging and electrophysiological experiments. The maximum ethanol

concentration used was relatively high, so particular attention was paid to these control experiments. In physiological saline, 0.5% ethanol did not produce a change in either the tone, the phasic contractions in any of the 12 preparations tested (number of animals,  $n_a = 9$ ), the WCTs (number of preparations,  $n_p = 9$ ,  $n_a = 3$ ), resting membrane potential (median  $-43.5$  mV; range  $-48.5$  to  $-32.7$  mV; to median  $-42.3$  mV; range  $-50.2$  to  $-31.9$  mV), or the amplitude of spontaneous depolarizations (median 4.2 mV; range 3.4 to 6.0 mV; to 4.0 mV; range 3.2 to 5.0 mV;  $P = \text{NS}$ ). The nomenclature of receptors and ion channels conforms to BJP's Guide to Receptors and Channels (Alexander *et al.*, 2011).

### Statistical analysis

The normality and homogeneity of variance were tested prior to further statistical analysis using Kolmogorov-Smirnov and Levene's tests, respectively (SPSS version 11; SPSS Inc., Chicago, IL, USA). Other statistical analyses were performed in Prism 4 (GraphPad Software, CA, USA). Unless otherwise stated, statistical comparisons were by Student's two-tailed paired *t*-tests. The term  $n_a$ , used in the presentation of statistical analyses throughout, refers to the number of animals used and the term  $n_p$  refers to the number of preparations used. To test for functional coupling between nearby cells, contingency tables were constructed and tested with Fisher's exact test (for a methodological example, see Brain *et al.*, 2002).


## Results

### Contraction studies

On mouse isolated bladder strips, PGE<sub>2</sub> (50  $\mu$ M) caused a concentration-dependent increase in tone with an EC<sub>50</sub> of 9.2  $\mu$ M (pEC<sub>50</sub> =  $5.04 \pm 0.41$ ; Figure 1A;  $n_p = 11$ ,  $n_a = 6$ ). The tone developed slowly (Figure 1B; median time to plateau: 56 s; range 41–160 s;  $n_p = 20$ ,  $n_a = 10$ ) and rapid phasic contractions were superimposed. The basal tone increased from  $2.26 \pm 0.33$  mN to  $4.51 \pm 0.32$  mN ( $P < 0.001$ ,  $n_p = 20$ ,  $n_a = 10$ ; Figure 1C). After stabilization of the tone, the rapid changes in tension continued until washout, although a reduction in frequency was observed 12–15 min after the application of PGE<sub>2</sub> (50  $\mu$ M) (by  $40.4 \pm 3.1\%$ ,  $n_p = 14$ ,  $n_a = 7$ ). Only five strips showed phasic contractions under control conditions and the average frequency was increased by PGE<sub>2</sub> (from  $0.042 \pm 0.016\text{ s}^{-1}$  to  $0.24 \pm 0.02\text{ s}^{-1}$ ;  $P < 0.01$ ,  $n_p = 15$ ,  $n_a = 8$ ; Figure 1D).

To determine whether part of this response could be driven by an action on efferent parasympathetic nerves, the effects of the Na<sup>+</sup> channel blocker TTX and the N-type Ca<sup>2+</sup> channel blocker  $\omega$ -conotoxin GVIA were investigated. TTX (1  $\mu$ M) did not affect the increase in basal tone observed after addition of PGE<sub>2</sub> (50  $\mu$ M) (relative change of  $5.3 \pm 5.5\%$  compared with PGE<sub>2</sub> alone;  $P = \text{NS}$ ,  $n_p = 8$ ,  $n_a = 6$ ), nor the frequency of phasic contractions ( $14 \pm 15\%$ ), nor their amplitude ( $14 \pm 9\%$ ). Similarly,  $\omega$ -conotoxin GVIA (100 nM) pre-treatment had no significant effect on PGE<sub>2</sub>-induced changes in basal tone ( $15 \pm 8\%$ ;  $P = \text{NS}$ ,  $n_p = 8$ ,  $n_a = 6$ ). Phasic contraction frequency ( $4.3 \pm 3.5\%$ ,  $P = \text{NS}$ ,  $n_p = 8$ ,  $n_a = 6$ ) and amplitude ( $9.1 \pm 7.8\%$ ,  $P = \text{NS}$ ,  $n_p = 8$ ,  $n_a = 6$ ) were also not significantly affected.





**Figure 1**

Exogenously applied PGE<sub>2</sub> contracts the murine urinary bladder strips in a concentration-dependent manner (A;  $n_p = 11$ ,  $n_a = 6$ ) with an EC<sub>50</sub> of 9.2 μM. (B) A sample trace of the effects of PGE<sub>2</sub> (50 μM) on such a muscle strip ( $n_p = 20$ ,  $n_a = 10$ ). The increase in tone is followed by phasic activity. PGE<sub>2</sub> (50 μM) increased tone (C;  $n_p = 20$ ,  $n_a = 10$ ) and the frequency of phasic contractions (D;  $n_p = 14$ ,  $n_a = 7$ ). \*\* $P < 0.01$  and \*\*\* $P < 0.001$ , compared to control.

To determine whether L-type Ca<sup>2+</sup> channels were required for the response to PGE<sub>2</sub> (50 μM), the selective L-type Ca<sup>2+</sup> channel blocker nifedipine (1 μM) was used. Nifedipine reduced the increase in basal tone observed after addition of PGE<sub>2</sub> (50 μM) by  $45.3 \pm 6.6\%$  ( $P < 0.05$ ,  $n_p = 20$ ,  $n_a = 10$ ) and abolished the PGE<sub>2</sub>-induced phasic contractions in all preparations (Figure 2Aa). To determine whether neurotransmitter release mediated PGE<sub>2</sub>-induced contractions, the contributions of purinergic and cholinergic neurotransmission were


explored by application of the P2X1 selective antagonist NF449 (10 μM) or the muscarinic receptor antagonist cyclopentolate (1 μM). The P2X1 selective antagonist NF449 (10 μM) did not significantly affect the PGE<sub>2</sub> (50 μM)-induced tone ( $8.4 \pm 12.4\%$ ,  $P = \text{NS}$ ,  $n_p = 15$ ,  $n_a = 9$ ), but significantly reduced the frequency of phasic contractions (by  $23.5 \pm 8.5\%$ ,  $P < 0.05$ ,  $n_p = 6$ ,  $n_a = 4$ ; Figure 2Ab). Cyclopentolate pretreatment had no effect on the PGE<sub>2</sub>-induced tone ( $15 \pm 18\%$ ,  $P = \text{NS}$ ,  $n_p = 8$ ,  $n_a = 4$ ) or on the frequency of phasic contractions ( $0.4 \pm 2.4\%$ ,  $P = \text{NS}$ ,  $n_p = 7$ ,  $n_a = 4$ ; Figure 2Ac).

In order to identify the role of EP receptors in the response to PGE<sub>2</sub> (50 μM), SC-51322 (10 μM), a selective EP<sub>1</sub> antagonist, and AH-6809 (10 μM), a non-selective EP<sub>2</sub> and EP<sub>3</sub> antagonist were used (Figure 2Ad). The binding affinity for SC-51322 is 14 nM for EP<sub>1</sub> receptors, and AH-6809 has a  $K_i$  of 1.1–1.6 μM at EP<sub>1–3</sub> receptors (Abramovitz *et al.*, 2000); so, both of these antagonists were used at concentrations significantly in excess of their  $K_i$  and/or binding affinity. When used together a significant reduction in the PGE<sub>2</sub> (50 μM)-induced tone was observed (by  $59.2 \pm 6.2\%$ ,  $P < 0.01$ ,  $n_p = 6$ ,  $n_a = 6$ ) and phasic contractions did not occur. When used separately SC-51322 and AH-6809 significantly reduced the PGE<sub>2</sub> (50 μM)-induced tone, but to a lesser extent (by  $39.1 \pm 5.8\%$ ,  $P < 0.05$ ,  $n_p = 7$ ,  $n_a = 6$  and  $23.7 \pm 4.4\%$ ,  $P < 0.05$ ,  $n_p = 5$ ,  $n_a = 5$ , respectively) suggesting an additive effect of the two antagonists. The frequency of phasic contractions was significantly reduced by SC-51322 ( $64 \pm 38\%$ ,  $P < 0.05$ ,  $n_p = 7$ ,  $n_a = 6$ ), but not by AH-6809 alone ( $9.7 \pm 4.7\%$ ,  $P = \text{NS}$ ,  $n_p = 5$ ,  $n_a = 5$ ). The effects of these antagonists on tone and the frequency of phasic contractions are summarized in Figure 2B and C, respectively.

### Electrophysiology

Electrophysiological studies were used to characterize the electrical changes in response to PGE<sub>2</sub> (50 μM). Spontaneous depolarizations, i.e. both spontaneous excitatory junction potentials (sEJPs) and sAPs, occurred with a stochastic temporal distribution in control conditions, but occurred in bursts following the superfusion of PGE<sub>2</sub> (50 μM; Figure 3). Bursting began following superfusion for 350 s (median; range 161 to 629 s;  $n_a = 11$ ). Bursts of spontaneous depolarizations occurred at a frequency of  $0.14 \pm 0.02$  Hz (mean  $\pm$  SEM; range 0.08 to 0.22 Hz, no. of bursts 10–138 from  $n_a = 10$ ). One way of showing that the spontaneous depolarizations occur in bursts is to measure the interval between successive events; if these events occur independently then the probability density function for such intervals should be exponential. This distribution of intervals between successive depolarizations in control conditions was exponential, but in the presence of PGE<sub>2</sub> events with a short (<2 s) interval occurred with a greater probability than expected (Figure 4A), implying that they were clustered in bursts.

PGE<sub>2</sub> (50 μM) increased the frequency of spontaneous depolarizations from 0.31 Hz (median; range 0.18 to 0.75 Hz) to 0.90 Hz (range 0.40 to 1.41 Hz; two-tailed paired  $t$ -test,  $P < 0.001$ ,  $n_p = 13$ ,  $n_a = 12$ ; Figure 4Ba). This is in contrast to the solvent (0.5% ethanol) control, which actually reduced the frequency of spontaneous depolarizations from 0.36 Hz (median; range 0.14 to 0.51 Hz) to 0.29 Hz (range 0.10 to 0.44 Hz) ( $P < 0.001$ ). PGE<sub>2</sub> (50 μM) also depolarized the resting membrane potential from  $-43.9$  mV (median; range



**Figure 2**

L-type  $\text{Ca}^{2+}$  channels and EP receptors mediate  $\text{PGE}_2$ -induced phasic contractions. (A) A comparison of the effects of the superfusion of the selective L-type  $\text{Ca}^{2+}$  channel blocker nifedipine (1  $\mu\text{M}$ ; a), the P2X1 receptor antagonist NF449 (10  $\mu\text{M}$ ; b), the muscarinic receptor antagonist cyclopentolate (1  $\mu\text{M}$ ; CP; c) and a combination of the selective EP<sub>1</sub> receptor antagonist SC-51322 (10  $\mu\text{M}$ ) and the non-selective EP<sub>2</sub> and EP<sub>3</sub> receptor antagonist AH-6809 (10  $\mu\text{M}$ ; d). The black trace shows a matching control trace (i.e. without any antagonist present), while the grey trace shows the response in the presence of the relevant antagonist(s). (B) A comparison of the change in tone for each drug application to normalized control frequencies; Student's paired *t*-test was used for the comparisons. The effect of each drug on the frequency of phasic activity was also assessed (C). \**P* < 0.05.

–50.2 to –24.6 mV) to –33.3 mV (range –41.2 to –16.1 mV) (*P* < 0.001; Figure 4Bb). In control conditions, depolarizations occurred with variable rise time (time from 10 to 90% peak amplitude) with a median of 18.8 ms (range 7.4 to 66.0 ms). Following superfusion of  $\text{PGE}_2$  (50  $\mu\text{M}$ ), the resulting spontaneous depolarizations occurred with a much broader rising phase, that is, with a greater rise time of 39.3 ms (range 20.8 to 98.5 ms) (*P* < 0.001; Figures 3C, 4Bc). The median amplitude of depolarizations increased from 4.0 mV (median; range 3.0 to 10.7 mV) in control conditions to 5.0 mV (4.5 to 16.7 mV) in the presence of  $\text{PGE}_2$  (50  $\mu\text{M}$ ) (*P* < 0.05; Figure 4Bd).

In order to test whether the effects of  $\text{PGE}_2$  were mediated through P2X1 receptors,  $\text{PGE}_2$  (50  $\mu\text{M}$ ) was superfused in combination with the selective P2X1 antagonist NF449 (10  $\mu\text{M}$ ). Pretreatment with NF449 (10  $\mu\text{M}$ ) did not change the amplitude, frequency or interval between spontaneous depolarizations, nor did it prevent the change in the resting membrane potential, in response to  $\text{PGE}_2$  (50  $\mu\text{M}$ ) (Figure 5A; Table 1). There was, however, an increase in the rise time of events from 47.7 ms with  $\text{PGE}_2$  (50  $\mu\text{M}$ ) alone to 67.4 ms when NF449 (10  $\mu\text{M}$ ) was also present (*P* < 0.05).

The L-type  $\text{Ca}^{2+}$  channel antagonist nifedipine (1  $\mu\text{M}$ ) prevented all measured effects of  $\text{PGE}_2$  on spontaneous


depolarizations, but the  $\text{PGE}_2$ -stimulated change in resting membrane potential remained (Figure 5B; Table 2).

### *Ca<sup>2+</sup> imaging*

$\text{Ca}^{2+}$ -imaging was used to determine the effect of  $\text{PGE}_2$  (50  $\mu\text{M}$ ) on smooth muscle  $\text{Ca}^{2+}$  regulation, particularly WCTs. Previous studies have shown that WCTs are due to smooth muscle action potentials (Heppner *et al.*, 2005; Young *et al.*, 2008).

Loading detrusor smooth muscle with the  $\text{Ca}^{2+}$  indicator Oregon Green BAPTA-1 AM and imaging groups of smooth muscle under control conditions revealed three types of spontaneous  $\text{Ca}^{2+}$  transients: WCTs,  $\text{Ca}^{2+}$  waves and local  $\text{Ca}^{2+}$  transients. Only WCT frequency and amplitude were investigated in this study.

WCTs were present under control conditions and their frequency was increased significantly by  $\text{PGE}_2$  (50  $\mu\text{M}$ ) (by  $72 \pm 5\%$ ; *P* < 0.001,  $n_p = 27$  cells,  $n_a = 9$ ; Figure 6A,B).  $\text{PGE}_2$  (50  $\mu\text{M}$ ) also appeared to increase the occurrence of subcellular  $\text{Ca}^{2+}$  transients, including waves (see Figure S1), although the frequency was not calculated. The amplitude of the WCTs was not affected by  $\text{PGE}_2$  ( $0.340 \pm 0.031 \Delta\text{F}/\text{F}_0$  under control conditions;  $0.300 \pm 0.025 \Delta\text{F}/\text{F}_0$  after  $\text{PGE}_2$  superfusion; *P* = NS; Figure 6C). The  $\text{Ca}^{2+}$  channel blocker nifedipine (1  $\mu\text{M}$ )




**Figure 3**

Superfusion of PGE<sub>2</sub> (50 μM) depolarizes SMCs and induces bursts of spontaneous depolarizations. (A) Over a period of 18 min, PGE<sub>2</sub> (50 μM) depolarized the resting membrane potential of a SMC within a strip of mouse isolated detrusor. (B) A comparison of the spontaneous activity prior to (Ba) and during (Bb) PGE<sub>2</sub> superfusion; PGE<sub>2</sub> stimulated the large amplitude spontaneous depolarizations that occur in bursts. (C) Shows the PGE<sub>2</sub>-stimulated spontaneous depolarizations have slow rise times.

prevented PGE<sub>2</sub>-induced WCTs (11 cells,  $n_a = 4$ ; Figure 7A). In the absence of PGE<sub>2</sub>, incubation with the selective P2X1 antagonist NF449 (10 μM) greatly reduced the frequency of WCTs (by  $98 \pm 6\%$ ;  $P < 0.001$ , two-way ANOVA,  $n_p = 18$  cells,  $n_a = 4$ ). Upon addition of PGE<sub>2</sub> (50 μM) in the presence of the NF449 (10 μM), WCTs were observed but at a significantly lower frequency than with PGE<sub>2</sub> alone (by  $28 \pm 17\%$ ;  $P < 0.001$ , two-way ANOVA; Figure 7B,C).

### PGE<sub>2</sub> and cellular coupling


PGE<sub>2</sub> (50 μM) superfusion increased the frequency of WCTs, but also nearby cells responded synchronously. To determine if such apparent synchronicity was random, contingency tables were used to compare the number of times



**Figure 4**

The distribution of intervals between successive spontaneous depolarizations (Aa) demonstrates that under control conditions they occur randomly, as the control distribution (solid line, black) follows the expected exponential distribution (dashed line, black). In the presence of PGE<sub>2</sub> (50 μM), bursting occurs, shown by the excess of brief intervals compared with the expected exponential distribution (compare the dashed and solid grey lines). (Ab) The median intervals were also briefer in the presence of PGE<sub>2</sub>. (B) PGE<sub>2</sub> also stimulated an increase in spontaneous depolarization frequency (a), an increase in rise time (c) and amplitude (d), and a depolarization of the RMP of SMCs (b). The data were obtained from  $n_p = 13$  and  $n_a = 12$  (Ab, B); \* $P < 0.05$  and \*\*\* $P < 0.001$ .

the cells responded separately against the number of times the cells responded together. In two of five pairs of cells tested there was a significant synchronous activity of WCTs after PGE<sub>2</sub> (50 μM) addition compared to no significant synchronous activity under control conditions for these same pairs ( $P < 0.001$ ; Fisher's exact test; see Figure 8A and Figure S2).



**Figure 5**  
PGE<sub>2</sub> (50 μM)-induced bursts of spontaneous depolarizations are mediated by L-type Ca<sup>2+</sup> channels and not P2X1 receptors. (A) The P2X1 receptor antagonist NF449 (1 μM) did not affect the occurrence of PGE<sub>2</sub> (50 μM)-induced bursts of spontaneous depolarizations. (B) The occurrence of these events was abolished by the L-type Ca<sup>2+</sup> channel antagonist nifedipine (1 μM).

To determine whether there was an effect of PGE<sub>2</sub> (50 μM) or its vehicle (0.5% ethanol) on electrical coupling between mouse detrusor SMCs, a series of brief intracellular current injections (−0.03 to −0.15 nA of 100 ms duration) was applied. Input resistances, calculated from the linear slope of the current–voltage relationship, varied greatly between SMCs. However, there was no significant difference between the input resistance of SMCs in control conditions, 212 MΩ (median; range 115 to 660 MΩ) and following superfusion of the vehicle (0.5% ethanol; 168 MΩ; range 84 to 769 MΩ; Wilcoxon matched two-tailed pairs test, *P* = NS, *n*<sub>a</sub> = 5). Similarly, PGE<sub>2</sub> (50 μM) did not change the input resistance (median control 181 MΩ; range 115 to 1060; PGE<sub>2</sub> median 209 MΩ; range 98 to 1045; Wilcoxon matched two-tailed pairs test, *P* = NS, *n*<sub>a</sub> = 6).

## Discussion and conclusions

In the urinary bladder, PGE<sub>2</sub> is implicated in physiological and pathophysiological conditions (Khan *et al.*, 1999; Schröder *et al.*, 2004; Masunaga *et al.*, 2006). Previous studies have shown that PGE<sub>2</sub> causes bladder overactivity when instilled intravesically (Schussler, 1990; Lee *et al.*, 2008) and causes contraction of isolated bladder tissue (Paea *et al.*, 1998). However, it is unclear whether PG's effects are neuronal or on the SMCs themselves, an issue the present study addresses.

PGE<sub>2</sub> (50 μM)-induced activity included an increased frequency of phasic contractions (85% increase), increased frequency of WCTs (72% increase) and a 66% increase in spontaneous depolarizations. To the authors' knowledge, this is the first study to show cellular electrical and Ca<sup>2+</sup> responses to PGE<sub>2</sub> in the bladder.

**Table 1**

Descriptive statistics of the effects of PGE<sub>2</sub> and the purinoceptor P2X1 antagonist NF449 (10 μM) on properties of SMCs and their spontaneous depolarizations (sDep)

Treatment		PGE <sub>2</sub> (50 μM)		PGE <sub>2</sub> (50 μM) + NF449 (10 μM)		Paired <i>t</i> -test, <i>P</i> PGE <sub>2</sub> + NF449 vs. control		PGE <sub>2</sub> + NF449 vs. PGE <sub>2</sub>	
Control		Control		Control		Control		Control	
RMP (mV)	−40.0 [−46.9 to −35.9]	−29.9 [−32.0 to −25.3]		−29.6 [−32.0 to −26.7]		<0.05		NS	
sDep									
Frequency (Hz)	0.5 [0.2–0.8]	0.6 [0.6–1.4]		0.7 [0.6–1.4]		<0.05		NS	
Amplitude (mV)	4.6 [3.6–8.6]	6.3 [4.6–13.9]		5.5 [4.0–10.9]		NS		NS	
Interval (s)	1.6 [1.1–3.8]	0.9 [0.6–1.3]		0.9 [0.6–1.1]		NS (0.07)		NS	
Rise time (ms)	19 [11–66]	48 [20–90]		67 [37–96]		<0.05		<0.05	
No. events	724 [304–951]	736 [453–888]		688 [286–1023]		–		–	
<i>n</i> <sub>a</sub>	4	4		4		4		4	

Data shown are median [range]. *n*<sub>a</sub>, number of animals. NS denotes *P* > 0.05.



**Table 2**

The effects of PGE<sub>2</sub> (50 µM) and the L-type Ca<sup>2+</sup> channel antagonist nifedipine (1 µM) on properties of SMCs and their spontaneous depolarizations (sDep)

	Control	PGE <sub>2</sub> (50 µM) + nifedipine (1 µM)	P
RMP (mV)	−43.8 [−60.7 to −36.5]	−33.5 [−55.8 to −24.9]	<0.01
sDep			
Frequency (Hz)	0.3 [0.1–0.7]	0.2 [0.0–0.6]	NS (0.29)
Amplitude (mV)	4.7 [3.3–5.4]	3.9 [3.4–5.4]	NS (0.08)
Interval (s)	2.6 [1.1–6.3]	2.9 [1.1–7.2]	NS (0.11)
Rise time (ms)	19 [12–25]	18 [9–21]	NS (0.29)
No. events	301 [148–2602]	569 [102–688]	
n <sub>a</sub>	7	7	

Data shown are median [range]. n<sub>a</sub>, number of animals. NS denotes  $P > 0.05$ .

Not all of the tissue strips displayed clear phasic contractions under control conditions, which may be due to different sensitivity to stretch, receptor subtype spatial inhomogeneity or other unidentified sources of variability amongst preparations.


### *PGE<sub>2</sub>-induced phasic contractions are mediated by L-type Ca<sup>2+</sup> channels and EP receptors*

Previous studies on mouse and rat urinary bladder have implicated both EP<sub>1</sub> and EP<sub>3</sub> receptors in PGE<sub>2</sub>-induced bladder overactivity. PF-2907617-02 and SC-19220, selective EP<sub>1</sub> antagonists, increased bladder capacity in normal rat bladders, but did not identify a necessary role of EP<sub>1</sub> receptors in normal micturition (Maggi *et al.*, 1988; Lee *et al.*, 2007). Thus it has been suggested that EP<sub>1</sub> receptors stimulate physiological effects during the storage phase but not during the voiding phase of micturition (Lee *et al.*, 2007). Cystometry on EP<sub>1</sub> KO mice supported the idea that EP<sub>1</sub> receptor function is not required for normal micturition (Maggi *et al.*, 1988). However, the PGE<sub>2</sub>-induced overactivity observed in rats was blocked by PF-2907617-02 in a dose-dependent manner (Lee *et al.*, 2007), and PGE<sub>2</sub>-induced overactivity in mice only occurred in wild-type (WT) EP<sub>1</sub> mice and not in KO EP<sub>1</sub> mice, implicating the EP<sub>1</sub> receptor directly with overactivity (Maggi *et al.*, 1988). In our experiments, blocking EP<sub>1</sub> receptors with SC-51322, which is effective in rodents (Moreland *et al.*, 2003), reduced the PGE<sub>2</sub>-induced frequency of phasic contractions (64% reduction) and decreased the change in tone observed after PGE<sub>2</sub> addition (39% reduction).

Other studies have implicated EP<sub>3</sub> receptors as mediators of PGE<sub>2</sub>-induced bladder overactivity. Infusion of PGE<sub>2</sub> into WT mice induced bladder overactivity, an effect that was not present in EP<sub>3</sub> KO mice. Furthermore, EP<sub>3</sub> KO mice had an enhanced bladder capacity compared to WT mice, therefore associating EP<sub>3</sub> receptors with normal bladder physiology (McCafferty *et al.*, 2008). In our experiments, non-selective EP<sub>2</sub> and EP<sub>3</sub> antagonism using AH-6809 caused a significant decrease in the change of tone induced by

PGE<sub>2</sub> (50 µM) ( $23.7 \pm 4.4\%$ ). However, AH-6809 did not significantly affect the frequency of phasic contractions observed after PGE<sub>2</sub> (50 µM) addition ( $9.7 \pm 4.7\%$ ). The effect of SC-51322 and AH-6809 (both 10 µM) were additive for an effect on tone after PGE<sub>2</sub> (50 µM) addition. The PGE<sub>2</sub>-induced increase in the frequency of phasic contractions was also abolished in all preparations in the presence of both antagonists. Thus, in order to reduce the PGE<sub>2</sub>-induced phasic contractions, blocking EP<sub>1</sub> receptors is necessary, but in order to prevent these phasic contractions all EP receptors present must be blocked. The change in tone could not be abolished with non-selective EP receptor antagonism, suggesting the involvement of another mechanism or that the antagonist concentration was insufficient to overcome an agonist concentration of about five times greater than the EC<sub>50</sub> (50 µM compared with 9.2 µM). Alternatively, degradation of PGE<sub>2</sub> by 15-hydroxyprostaglandin dehydrogenase, which is known to be expressed by urothelial cells (Tseng-Rogenski *et al.*, 2008), might initiate a catabolic cascade of which some members may be active.

In our studies, we found that blocking L-type Ca<sup>2+</sup> channels with nifedipine abolished all PGE<sub>2</sub>-induced phasic contractions, WCTs and sAPs. In addition, it reduced the change in tone observed after PGE<sub>2</sub> addition by  $45.3 \pm 6.6\%$ . This indicates that PGE<sub>2</sub>-induced phasic contractions require smooth muscle L-type Ca<sup>2+</sup> channels and EP receptors on smooth muscle or other nearby cells. Previous studies using the L-type Ca<sup>2+</sup> channel antagonist verapamil showed similar effects on contraction (Morita *et al.*, 1994); together, these findings suggest that PGE<sub>2</sub> acts to increase Ca<sup>2+</sup> influx into the cell. There is evidence that the signal transduction of EP<sub>1</sub> involves extracellular Ca<sup>2+</sup> entry through a pathway independent of phospholipase C activation (Katoh *et al.*, 1995). Given the PGE<sub>2</sub>-induced depolarization we now report, it seems that this slow depolarization causes bursts of L-type Ca<sup>2+</sup> channel-dependent smooth muscle action potentials. The remaining questions are how such a slow depolarization is initiated by PGE<sub>2</sub> and how the residual contraction observed on blocking L-type Ca<sup>2+</sup> channels is mediated.




**Figure 6**

PGE<sub>2</sub> increases the frequency of WCTs, but not their amplitude. (A) A sample trace of the frequency of WCTs. This trace is a concatenated composite of shorter 19.8 s recording segments taken every minute. The scale bar indicates the time over the whole experiment so is not valid within each segment. The trace shows the presence of WCTs under control conditions and after addition of PGE<sub>2</sub> (50 μM). The positive-going spikes on the trace are WCTs, shown throughout the course of an experiment for a typical preparation; slow changes in the baseline represent focal plane drift while imaging and are hence an artefact of this non-ratioable Ca<sup>2+</sup> indicator. (B) The effect of PGE<sub>2</sub> (50 μM) on WCT frequency. (C) The effect of PGE<sub>2</sub> (50 μM) on WCT amplitude. \*\**P* < 0.01.


### *PGE<sub>2</sub>-induced phasic contractions are not neurogenic*

There is evidence for EP<sub>3</sub> receptors on some cholinergic parasympathetic nerve terminals (Reinheimer *et al.*, 1998; Spicuzza *et al.*, 1998), facilitation of parasympathetic neurotransmitter release through EP<sub>1</sub> receptors (Jadhav *et al.*, 2009) and direct depolarization of intrinsic cardiac neurons by PGE<sub>2</sub> through SK<sub>Ca</sub> channel inhibition (Jelson *et al.*, 2003). All of these findings led to the suggestion that PGE<sub>2</sub> induces similar facilitatory effects in the urinary bladder. However, none of the observed effects of PGE<sub>2</sub> in the mouse bladder were prevented by TTX, which is known to abolish field-stimulation-induced contraction in the mouse bladder (Holt *et al.*, 1985), indicating that action potential generation in efferent nerves was not required. This observation is consis-



**Figure 7**

PGE<sub>2</sub>-induced increase in frequency of WCTs is abolished by nifedipine and reduced by NF449. (A) WCTs after the addition of PGE<sub>2</sub> (50 μM) alone and in the presence of nifedipine (1 μM). The spikes on the trace represent WCTs, against time, in one preparation. (B) The frequency of WCTs after superfusion with PGE<sub>2</sub> (50 μM) in the presence and absence of NF449 (10 μM). Each trace is a concatenated composite of shorter 13.7 s recording segments taken every minute. (C) A comparison of the frequency of WCTs under different conditions. \**P* < 0.05 and \*\**P* < 0.01.



**Figure 8**

WCTs occur simultaneously across multiple neighbouring smooth muscle cells in the presence of PGE<sub>2</sub>. (A) A region of a mouse isolated bladder detrusor (loaded with the Ca<sup>2+</sup> indicator Oregon Green 488 BAPTA-1 AM) showing four smooth muscle cells (denoted by asterisks). Time frames are 198 ms apart, arranged horizontally. The total number of frames sampled was 350. All cells responded to PGE<sub>2</sub> and the cells labelled a and b were analysed to test if the probability of a WCT occurring synchronously in both cells was increased after PGE<sub>2</sub> addition. (B) The frequency of synchronous and asynchronous WCTs under control conditions and after the addition of PGE<sub>2</sub>. \**P* < 0.05 (contingency tables and Fisher's exact test). Only in the presence of PGE<sub>2</sub> were synchronous WCTs detected at a rate greater than that predicted by chance.

tent with experiments in human bladder strips, demonstrating that the response to PGE<sub>2</sub> is insensitive to TTX (Andersson *et al.*, 1977). Additionally, the failure of ω-conotoxin GVIA to inhibit the response also contravenes the idea that local depolarization of the nerve terminals triggers local nerve terminal Ca<sup>2+</sup> influx and transmitter release, although it is acknowledged that some voltage-gated Ca<sup>2+</sup> channel subtypes

other than N-type Ca<sup>2+</sup> channels can contribute to parasympathetic transmission in the mouse bladder (Waterman, 1996).

In the normal mouse bladder spontaneous depolarizations and sAPs result from nerve-released ATP acting on P2X1 receptors (Meng *et al.*, 2008; Young *et al.*, 2008), although under other experimental conditions there may also be spontaneous activity that is purely myogenic (Hayase *et al.*, 2009). The first biophysical indication that PGE<sub>2</sub>-induced depolarizations are induced in a different manner to the normal sEJPs and sAPs was the difference in their electrical properties, such as a slower rise time (Table 1). We found that NF449 alone abolished sEJPs, sAPs and WCTs (confirming the earlier observations of our group; Young *et al.*, 2008), yet PGE<sub>2</sub> was still able to initiate similar events, albeit with a lower frequency of WCTs (by  $28 \pm 17\%$ ) and phasic contractions (by  $23.5 \pm 8.5\%$ ). Interestingly, the reduction in the frequency of PGE<sub>2</sub>-induced WCTs observed in the presence of NF449 (to  $0.050 \pm 0.009$  Hz) is similar to the control frequency of WCTs ( $0.053 \pm 0.027$  Hz). While PGE<sub>2</sub>-induced spontaneous depolarizations were not affected by NF449 (Table 2), we feel that this represents a lack of statistical power (*n* = 4 cells) in this technically demanding set of experiments. Also, PGE<sub>2</sub>-induced phasic contractions in the presence of NF449 were reduced (to  $0.103 \pm 0.065$  Hz) to a level similar to the control frequency of phasic contractions ( $0.062 \pm 0.044$  Hz). Overall, these data suggest that pretreatment with NF449 reduces basal neurogenic, purinergic activity but not the PGE<sub>2</sub>-induced activity. PGE<sub>2</sub>-induced phasic contractions were insensitive to the muscarinic antagonist cyclopentolate (1 μM). These results argue against both direct and indirect effects of PGE<sub>2</sub> on parasympathetic nerves innervating the urinary bladder, suggesting a direct action on the smooth muscle or other nearby cells.

### PGE<sub>2</sub> induces bursts of spontaneous activity

The synchronicity of WCTs in the presence of PGE<sub>2</sub> led to the investigation of whether PGE<sub>2</sub> increased electrical coupling between the cells. However, no significant fall in the input resistance (which represents a mixture of membrane resistance and gap junction resistance) was observed. It seems likely that the SMCs become functionally coupled not because of changes in gap junction resistance, but because of their depolarization. As cells depolarize (in this case by 11 mV), they move closer to their threshold for action potential initiation and hence require less depolarizing current to flow from adjacent active cells in order to initiate and propagate action potentials. In this way, depolarization directly increases functional action potential coupling, and hence co-ordinates micromotion or phasic contractions, without any change in the electrical resistance between cells.

K<sup>+</sup> channel inhibition may be responsible for the depolarization and the shift from stochastic action potential occurrence to bursting observed in the current study, similar to that observed in the guinea-pig bladder when small conductance Ca<sup>2+</sup>-activated K<sup>+</sup> (SK) channels are inhibited with apamin (Hashitani *et al.*, 2004a). In the mouse urinary bladder, apamin can increase the frequency of spontaneous depolarizations without changing the resting membrane potential (Hayase *et al.*, 2009), while suppression of SK channels in mice increases the frequency of non-voiding contrac-

tions (Herrera *et al.*, 2003).  $K_v$  channels might also be involved; tetraethylammonium (TEA; 10 mM) acutely depolarizes mouse bladder SMCs (Hayase *et al.*, 2009), suppresses after-hyperpolarizations, and increases sAP frequency, although this effect was lost over time at that high concentration of TEA. WCTs also became more synchronous in TEA. So, although several  $K^+$  channels might be involved, inhibition of the TEA-sensitive (i.e.  $K_v$ ) channel most readily explains the effects of  $PGE_2$ , although this was not directly tested.

Chloride channel activation by  $PGE_2$  cannot be ruled out, because  $PGE_2$ -induced  $Cl^-$  channel activation has been shown in some other cell types (Okamoto *et al.*, 2004; Seto *et al.*, 2008). However, the driving force for  $Cl^-$  is much smaller than that for  $K^+$ , so a much larger change in  $Cl^-$  conductance would be needed in order to explain any given depolarization. Given that a significant change in input resistance was not detected (median change from 181 to 209 M $\Omega$ ), it seems unlikely that such a large fall in membrane resistance occurred.

### Interstitial cells and the role of PGs

Interstitial cells of Cajal (ICCs) are of great contemporary interest in urogenital physiology because of the close association of the cells with SMCs and nerves, and because of the presence of a large suburothelial plexus in the lamina propria (Brading and McCloskey, 2005). During the present study we were unable to identify cells with a morphology typical of ICCs, which responded to PGs with a change in  $Ca^{2+}$ , although quiescent cells with this morphology were observed in some series (see Figure S3 for an example). The experimental protocol was not optimized to investigate such cells, in part because viewing from the serosal side means that the lamina propria layer of ICCs (ICC-LP) cannot be seen. The more scattered detrusor ICCs, typically located at the edge of muscle bundles (see McCloskey, 2010 for a review) are difficult to differentiate morphologically from both Schwann cells and fibroblasts, and this study was adequately designed appropriately to study these cells. A vital staining method for ICCs would greatly facilitate such experiments. Thus, it is possible that some of the effects of  $PGE_2$  could have been through an indirect effect on another cell type (such as ICCs, urothelial cells or Schwann cells) releasing a paracrine factor, or coupling through gap junctions, which influenced SMCs (rather than the nerves). However, while there is strong evidence of active calcium transients and electrical activity in ICCs, these are not temporally coupled with either contraction of the smooth muscle, smooth muscle  $Ca^{2+}$  activity or smooth muscle electrical activity (Hashitani *et al.*, 2004b). Therefore, at the current time the involvement of ICCs is speculative.

ICCs may also act as a source of  $PGE_2$ , as they are known to release PGs in the urogenital tract under some conditions (Hashitani, 2006). This may mediate urothelial to SMC signalling.

### Implications of this study

$PGE_2$  is thought to play a role in bladder overactivity. *In situ*, the bladder's afferent nerves can stimulate spinal cord-mediated reflexes through C-fibre afferent nerves; PGs could

act as neuromodulators to induce this excitability and bladder overactivity, for example in the case of spinal cord injury (Maggi *et al.*, 1988).

Our experiments suggest that in addition to this reflex pathway there is a direct local action of  $PGE_2$  in the bladder that does not require an increase in cholinergic or purinergic parasympathetic activation. It is known that EP<sub>1</sub> KO mice do not develop bladder hyperactivity after bladder outlet obstruction but that WT mice do (Schröder *et al.*, 2004), highlighting the importance of prostanoid signalling for the pathogenesis of bladder overactivity.

Thus, we confirmed that  $PGE_2$  (50  $\mu$ M) increases smooth muscle activity in mouse bladder strips *in vitro*, as seen in other species (Maggi *et al.*, 1988). This was seen as an increase in phasic contraction, mediated by SMC depolarization and L-type  $Ca^{2+}$  channel-dependent action potentials (observed with both electrophysiology and  $Ca^{2+}$  imaging). Our studies also demonstrated that more than one EP receptor subtype can generate such spontaneous activity. Depolarization, rather than a fall in gap junction resistance, increases the functional coupling of SMCs. While the effects of prostanoids on afferent C-fibres are undoubtedly important, the direct effects of  $PGE_2$  on the isolated bladder (which do not require neurotransmitter release from the efferent parasympathetic nerves) implies that rational therapies for such overactivity should not focus solely on the spinal cord reflex arc, but should also prevent PG's direct action on bladder smooth muscle. These muscle-mediated effects could produce the detrusor overactivity that follows outlet obstruction and spinal cord injury, which are both associated with increases in  $PGE_2$  production (Masick *et al.*, 2001; Masunaga *et al.*, 2006).

### Acknowledgements

This work is supported by a Wellcome Trust Career Development Research Fellowship (074128 to KLB); a Wellcome Trust VIP award supported JSY, who is currently funded by an Age UK research fellowship.

### Conflicts of interest

None.

### References

- Abramovitz M, Adam M, Boie Y, Carriere M, Denis D, Godbout C *et al.* (2000). The utilization of recombinant prostanoid receptors to determine the affinities and selectivities of prostaglandins and related analogs. *Biochim Biophys Acta* 1483: 285–293.
- Alexander SPH, Mathie A, Peters JA (2011). Guide to Receptors and Channels (GRAC), 5th edition. *Br J Pharmacol* 164: S1–S324.
- Andersson KE, Arner A (2004). Urinary bladder contraction and relaxation: physiology and pathophysiology. *Physiol Rev* 84: 935–986.



- Andersson KE, Ek A, Persson CG (1977). Effects of prostaglandins on the isolated human bladder and urethra. *Acta Physiol Scand* 100: 165–171.
- Brading AF, McCloskey KD (2005). Mechanisms of Disease: specialized interstitial cells of the urinary tract – an assessment of current knowledge. *Nat Clin Pract Urol* 2: 546–554.
- Brain KL, Jackson VM, Trout SJ, Cunnane TC (2002). Intermittent ATP release from nerve terminals elicits focal smooth muscle  $\text{Ca}^{2+}$  transients in mouse vas deferens. *J Physiol* 541: 849–862.
- Bultitude MI, Hills NH, Shuttleworth KE (1976). Clinical and experimental studies on the action of prostaglandins and their synthesis inhibitors on detrusor muscle in vitro and in vivo. *Br J Urol* 48: 631–637.
- Cardozo LD, Stanton SL (1980). A comparison between bromocriptine and indomethacin in the treatment of detrusor instability. *J Urol* 123: 399–401.
- Dixon JS, Gosling JA (1990). Ultrastructure of smooth muscle cells in the urinary system. In: Motta PM (ed.). *Ultrastructure of Smooth Muscle*. London: Kluwer Academic, pp. 153–169.
- Dveksler G, Franchi AM, Gonzalez ET, Gimeno MA, Gimeno AL (1989). Electric field stimulation alters the outputs of prostaglandins from isolated rat urinary bladder preparations. Influences of papaverine and tetrodotoxin. *Prostaglandins Leukot Essent Fatty Acids* 36: 65–68.
- de Groat WC, Yoshimura N (2001). Pharmacology of the lower urinary tract. *Annu Rev Pharmacol Toxicol* 41: 691–721.
- Hanawa A (1991). Roles of prostaglandin on rabbit vesicourethral smooth muscle contraction. *Nippon Hinyokika Gakkai Zasshi* 82: 1256–1264.
- Hashitani H (2006). Interaction between interstitial cells and smooth muscles in the lower urinary tract and penis. *J Physiol* 576: 707–714.
- Hashitani H, Brading AF, Suzuki H (2004a). Correlation between spontaneous electrical, calcium and mechanical activity in detrusor smooth muscle of the guinea-pig bladder. *Br J Pharmacol* 141: 183–193.
- Hashitani H, Yanai Y, Suzuki H (2004b). Role of interstitial cells and gap junctions in the transmission of spontaneous  $\text{Ca}^{2+}$  signals in detrusor smooth muscles of the guinea-pig urinary bladder. *J Physiol* 559: 567–581.
- Hayase M, Hashitani H, Kohri K, Suzuki H (2009). Role of  $\text{K}^{+}$  channels in regulating spontaneous activity in detrusor smooth muscle in situ in the mouse bladder. *J Urol* 181: 2355–2365.
- Heppner TJ, Bonev AD, Nelson MT (2005). Elementary purinergic  $\text{Ca}^{2+}$  transients evoked by nerve stimulation in rat urinary bladder smooth muscle. *J Physiol* 564: 201–212.
- Herrera GM, Pozo MJ, Zvara P, Petkov GV, Bond CT, Adelman JP *et al.* (2003). Urinary bladder instability induced by selective suppression of the murine small conductance calcium-activated potassium (SK3) channel. *J Physiol* 551: 893–903.
- Holt SE, Cooper M, Wyllie JH (1985). Evidence for purinergic transmission in mouse bladder and for modulation of responses to electrical stimulation by 5-hydroxytryptamine. *Eur J Pharmacol* 116: 105–111.
- Ishizuka O, Mattiasson A, Andersson KE (1995). Prostaglandin  $\text{E}_2$ -induced bladder hyperactivity in normal, conscious rats: involvement of tachykinins? *J Urol* 153: 2034–2038.
- Jadhav V, Jabre A, Chen MF, Lee TJ (2009). Presynaptic prostaglandin  $\text{E}_2$   $\text{EP}_1$ -receptor facilitation of cerebral nitroergic neurogenic vasodilation. *Stroke* 40: 261–269.
- Jelson GS, DeMasi GM, Sager KL, Hardwick JC (2003). Modulation of guinea pig intrinsic cardiac neurons by prostaglandins. *Am J Physiol Regul Integr Comp Physiol* 285: R682–R689.
- Katoh H, Watabe A, Sugimoto Y, Ichikawa A, Negishi M (1995). Characterization of the signal transduction of prostaglandin  $\text{E}$  receptor  $\text{EP}_1$  subtype in cDNA-transfected Chinese hamster ovary cells. *Biochim Biophys Acta* 1244: 41–48.
- Khan MA, Thompson CS, Mumtaz FH, Jeremy JY, Morgan RJ, Mikhailidis DP (1998). Role of prostaglandins in the urinary bladder: an update. *Prostaglandins Leukot Essent Fatty Acids* 59: 415–422.
- Khan MA, Thompson CS, Angelini GD, Morgan RJ, Mikhailidis DP, Jeremy JY (1999). Prostaglandins and cyclic nucleotides in the urinary bladder of a rabbit model of partial bladder outlet obstruction. *Prostaglandins Leukot Essent Fatty Acids* 61: 307–314.
- Lee T, Hedlund P, Newgreen D, Andersson KE (2007). Urodynamic effects of a novel  $\text{EP}_1$  receptor antagonist in normal rats and rats with bladder outlet obstruction. *J Urol* 177: 1562–1567.
- Lee T, Andersson KE, Streng T, Hedlund P (2008). Simultaneous registration of intraabdominal and intravesical pressures during cystometry in conscious rats – effects of bladder outlet obstruction and intravesical  $\text{PGE}_2$ . *Neurourol Urodyn* 27: 88–95.
- Maggi CA (1992). Prostanoids as local modulators of reflex micturition. *Pharmacol Res* 25: 13–20.
- Maggi CA, Giuliani S, Patacchini R, Conte B, Furio M, Santicioli P *et al.* (1988). The effect of SC-19220, a prostaglandin antagonist, on the micturition reflex in rats. *Eur J Pharmacol* 152: 273–279.
- Masick JM, Levin RM, Hass MA (2001). The effect of partial outlet obstruction on prostaglandin generation in the rabbit urinary bladder. *Prostaglandins Other Lipid Mediat* 66: 211–219.
- Masunaga K, Yoshida M, Inadome A, Iwashita H, Miyamae K, Ueda S (2006). Prostaglandin  $\text{E}_2$  release from isolated bladder strips in rats with spinal cord injury. *Int J Urol* 13: 271–276.
- McCafferty GP, Misajet BA, Laping NJ, Edwards RM, Thorneloe KS (2008). Enhanced bladder capacity and reduced prostaglandin  $\text{E}_2$ -mediated bladder hyperactivity in  $\text{EP}_3$  receptor knockout mice. *Am J Physiol Renal Physiol* 295: F507–F514.
- McCloskey KD (2010). Interstitial cells in the urinary bladder – localization and function. *Neurourol Urodyn* 29: 82–87.
- Meng E, Young JS, Brading AF (2008). Spontaneous activity of mouse detrusor smooth muscle and the effects of the urothelium. *Neurourol Urodyn* 27: 79–87.
- Moreland RB, Kim N, Nehra A, Goldstein I, Traish A (2003). Functional prostaglandin  $\text{E}$  ( $\text{EP}$ ) receptors in human penile corpus cavernosum. *Int J Impot Res* 15: 362–368.
- Morita T, Ando M, Kihara K, Kitahara S, Ishizaka K, Matsumura T *et al.* (1994). Effects of prostaglandins  $\text{E}_1$ ,  $\text{E}_2$  and  $\text{F}_2$  alpha on contractility and cAMP and cGMP contents in lower urinary tract smooth muscle. *Urol Int* 52: 200–203.
- Okamoto F, Kajiya H, Fukushima H, Jimi E, Okabe K (2004). Prostaglandin  $\text{E}_2$  activates outwardly rectifying  $\text{Cl}^-$  channels via a cAMP-dependent pathway and reduces cell motility in rat osteoclasts. *Am J Physiol Cell Physiol* 287: C114–C124.
- Palea S, Toson G, Pietra C, Trist DG, Artibani W, Romano O *et al.* (1998). Pharmacological characterization of thromboxane and prostanoid receptors in human isolated urinary bladder. *Br J Pharmacol* 124: 865–872.



- Reinheimer T, Harnack E, Racke K, Wessler I (1998). Prostanoid receptors of the EP<sub>3</sub> subtype mediate inhibition of evoked [<sup>3</sup>H]acetylcholine release from isolated human bronchi. *Br J Pharmacol* 125: 271–276.
- Schröder A, Newgreen D, Andersson KE (2004). Detrusor responses to prostaglandin E<sub>2</sub> and bladder outlet obstruction in wild-type and Ep<sub>1</sub> receptor knockout mice. *J Urol* 172: 1166–1170.
- Schussler B (1990). Comparison of the mode of action of prostaglandin E<sub>2</sub> (PGE<sub>2</sub>) and sulprostone, a PGE<sub>2</sub>-derivative, on the lower urinary tract in healthy women. A urodynamic study. *Urol Res* 18: 349–352.
- Seto V, Hirota C, Hirota S, Janssen LJ (2008). E-ring isoprostanes stimulate a Cl<sup>−</sup> conductance in airway epithelium via prostaglandin E<sub>2</sub>-selective prostanoid receptors. *Am J Respir Cell Mol Biol* 38: 88–94.
- Spicuzza L, Giembycz MA, Barnes PJ, Belvisi MG (1998). Prostaglandin E<sub>2</sub> suppression of acetylcholine release from parasympathetic nerves innervating guinea-pig trachea by interacting with prostanoid receptors of the EP<sub>3</sub>-subtype. *Br J Pharmacol* 123: 1246–1252.
- Tseng-Rogenski S, Lee IL, Gebhardt D, Fischer SM, Wood C, Park JM *et al.* (2008). Loss of 15-hydroxyprostaglandin dehydrogenase expression disrupts urothelial differentiation. *Urology* 71: 346–350.
- Waterman SA (1996). Multiple subtypes of voltage-gated calcium channel mediate transmitter release from parasympathetic neurons in the mouse bladder. *J Neurosci* 16: 4155–4161.
- Young JS, Meng E, Cunnane TC, Brain KL (2008). Spontaneous purinergic neurotransmission in the mouse urinary bladder. *J Physiol* 586: 5743–5755.

## Supporting information

Additional Supporting Information may be found in the online version of this article:

**Figure S1** In addition to the effects on WCTs, PGE<sub>2</sub> appeared to increase the frequency of subcellular Ca<sup>2+</sup> transients, including waves. Some of these subcellular Ca<sup>2+</sup> transients also appear to be periodic. A control series of recordings and a subsequent recording from the same field of view in the presence of PGE<sub>2</sub> (50 μM) are shown. The scale bar represents 5 μm; for display purposes this movie plays at twice the sampling rate.

**Figure S2** PGE<sub>2</sub>-induced coupled WCTs in smooth muscle cells. A series of confocal images showing WCTs temporally synchronized in neighbouring cells – those shown in Figure 8A. Each frame is 50 μm across; for display purposes this movie plays at twice the sampling rate.

**Figure S3** Labelling of spindle-shaped cells reminiscent of ICCs. In most experiments, only smooth muscle (sm) cells were identified; in some experimental series, spindle-shaped cells reminiscent of ICCs, Schwann cells or fibroblasts (IC?) could be identified, brightly labelled with the Ca<sup>2+</sup> indicator. To date, such cells have been unresponsive to physiological stimuli. The field size is 100 μm.

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.